

QA
531
B6

UC-NRLF






\$B 320 203


YB 56842

IN MEMORIAM

Edward Bright







ACADEMIC

TRIGONOMETRY.

PLANE AND SPHERICAL.

BY

T. M. BLAKSLEE, PH.D. (YALE),
PROFESSOR OF MATHEMATICS IN THE UNIVERSITY OF DES MOINES.



BOSTON:
PUBLISHED BY GINN & COMPANY.
1888.

QA531
B6

*J. M.
Edward Bright*

Entered, according to Act of Congress, in the year 1887, by

T. M. BLAKSLEE,

in the Office of the Librarian of Congress, at Washington.

TYPOGRAPHY BY J. S. CUSHING & Co., BOSTON.

PRESSWORK BY GINN & Co., BOSTON.

BC

PREFACE.

THE purpose of this arrangement is to aid the memory by noting *analogies*.

$+$ and a^2 have as *spherical analogies* \times and $\cos a$. Page 12 has page 13, and each "Law" has its analogy.

In Spherical Trigonometry we note the *determining groups*: side, $+$, co. function, and \angle , $-$, function.

It is hoped that the Introduction will fix the characteristics of Trigonometry.

This should be accompanied by practical work, and occupy at least a week.

T. M. BLAKSLEE.

DES MOINES, IA.

NOTE. It is convenient for examinations to have tables separate from formulas.

The explanation of the use of tables should be with them.

Two pages of model solutions and answers may be added. (Opinions asked on this point.)

Digitized by the Internet Archive
in 2008 with funding from
Microsoft Corporation

INTRODUCTION.

DEF. Trigonometry is, etymologically, the Science of Measuring Triangles. Besides this, it now includes the Science of Angular Functions.


We first inquire, What is a function? then, What are the angular functions?

A function of a variable is a second variable so related to the first that any change in the variable produces a change in the function.

ILL. Oil in lamp and time it has burnt.

DEF. The functions of the angle between two straight lines are *the six ratios of the sides of the right triangle* formed by these lines and a perpendicular upon one of them from a point in the other.

NOTATION. h , hypotenuse; o , opposite; a , adjacent.



The ratios are, by definition,

$$\text{sine} = \sin = \frac{o}{h} \quad \therefore o = h \sin, \quad h = \frac{o}{\sin}$$

$$\text{cosine} = \cos = \frac{a}{h} \quad \therefore a = h \cos, \quad h = \frac{a}{\cos}$$

$$\text{tangent} = \tan = \frac{o}{a} \quad \therefore o = a \tan, \quad a = \frac{o}{\tan}$$

And their reciprocals,

$$\frac{\sin}{\cos} = \frac{o}{h} \div \frac{a}{h} = \frac{o}{a} = \tan.$$

By similar triangles, the functions are constants for a constant angle, but variables for a variable angle.

The base line is the initial line; the hypotenuse line, the terminus of the angle.

Linear Representation. (1) If $h = 1$, $o = \sin$, $a = \cos$.
(2) If $a = 1$, $o = \tan$.

The *transverse line* is TT' through vertex and perpendicular to initial line II' .

\sin = transverse projection of directed unit. (Unit h .)

\cos = initial projection of directed unit.*


\tan = transverse projection of h if initial projection be unity.†

Since antecedent = consequent \times ratio, also

For sine and cosine, consequent = h , ratio = function.

RULE I. To obtain either side from h , multiply by ratio, sine for o , cosine for a .

RULE II. To obtain the sine from cosine, multiply by tangent.



Quadrants. II' and TT' divide the angular space about the vertex into four quadrants, numbered as in the figure.

An angle is in the quadrant in which it terminates.

* The angle being the direction of its terminus, we may speak of the ratios as *direction ratios*.

Since for the other acute angle of ratio triangle,

$$\sin = \frac{a}{h}, \cos = \frac{o}{h}, \text{ and } \tan = \frac{a}{o} = \cotangent = \cot.$$

\therefore "co" means of complement.

† If a circle be described with the unit base a as radius, o is a tangent.

The Terminal Values of the functions are as follows :

	I.	II.	III.	IV.	∠	0°.	90°.	180°.	270°.
sin	+	+	—	—	sin	0	+1	0	—1
cos	+	—	—	+	cos	+1	0	—1	0
tan	+	—	+	—	tan	0	∞	0	∞


The algebraic signs being determined thus : to right and up, + ; to left and down, —.

PRACTICAL DEVELOPMENT.

Wishing to calculate the distance IB to an object B , starting from I , I laid off $IA \perp IB$.

At a distance $AM = 1$ from A I erected $MN \perp AI$, determining N by looking from A to B .

I also measured AP , and drew $PL \perp AB$.




The last is not needed in measuring the distance ; in fact, AM might have been any distance, when IB could have been found, as $IB = \frac{MN \times AI}{AM}$.

The advantage of a table of tangents is, that we never have need to construct the small triangle.

If $IA = 1000$ feet, and we have the tangent from a table, we have simply to move the decimal point three places, and we have IB at once.

Two-Place Table. Take 10 inches as an hypotenuse, and, by aid of a protractor (or by constructing an angle of 30° , geometrically, and then *trisecting it by folding*), construct the

values of sine and cosine ($\therefore \tan = \frac{\sin}{\cos}$) for every 10° . Here 10 inches = unit. \therefore 0.1 inch = 0.01 unit.



Evidently (arithmetically) function $(180^\circ - A) = f(A)$. The ratio triangles being equal, having h and A equal.

EXAMPLES.

\angle°	10	20	30	40	50	60	70	80
sin	17	34	50	64	77	87	94	98
cos	98	94	87	77	64	50	34	17
tan	18	36	58	84	1.19	1.73	2.75	5.67

1. Give functions, if o , a , h , are (1) 6, 8, 10; (2) 10, 24, 26; (3) 4, 7, 5, 8.5.

2. Solve the following: \angle , o , a , h being (1) 20° , ?, ?, 100; (2) ?, 4, ?, 5; (3) $57'$, 4000, ?, ?; (4) $8.8''$, 4000, ?, ?.

NOTE. If the greatest angular distance of Venus from the sun be 45° , what is its distance from that body as compared to that of the earth?

3. Can the sines of 0° , 30° , 45° , 60° , and 90° be written, $\frac{1}{2}\sqrt{0}$, $\frac{1}{2}\sqrt{1}$, $\frac{1}{2}\sqrt{2}$, $\frac{1}{2}\sqrt{3}$, $\frac{1}{2}\sqrt{4}$?

4. If A , B , and C be the angles, and a , b , and c the opposite sides of a triangle, p the perpendicular from C to c , show that $a \sin B = p = b \sin A$. $\therefore a : b = \sin A : \sin B$. (In words.)

Do field work, using ratios to two places,

$$\sin 15 = \frac{1}{2}(0.17 + 0.34),$$

$$\sin 18 = \sin 10 + 0.8 \times 0.17 = 0.31.$$

NOTE. If the greatest and least values of the maximum elongation of Mercury be 15° and 30° , what are its greatest and least distances from the sun?

Logarithmic Solutions. Though strictly Algebra, we give the logarithmic solutions thus far:

$$\log \text{ of } \sin = \log o - \log h. \quad \therefore \log o = \log h + \log \text{ of } \sin.$$


$$\log \text{ of } \cos = \log a - \log h. \quad \therefore \log a = \log h + \log \text{ of } \cos.$$

$$\log \text{ of } \tan = \log o - \log a. \quad \therefore \log o = \log a + \log \text{ of } \tan.$$

$$\log \sin = \log \text{ of } \sin + 10, \quad a : b = \sin A : \sin B, \text{ gives}$$

$$\log \sin A = \log a + \text{colog } b + \log \sin B.$$

PLANE.



THE DEFINING EQUATIONS.

$$(1) \text{ By } T_1, \sin A = \frac{a}{h}, \quad a = h \sin A, \quad h = \frac{a}{\sin A}.$$

$$(2) \text{ By } T_3, \cos A = \frac{b}{h}, \quad b = h \cos A, \quad h = \frac{b}{\cos A}.$$

$$(3) \text{ By } T_2, \tan A = \frac{a}{b}, \quad a = b \tan A, \quad b = \frac{a}{\tan A}.$$

(1) is the definition of the *sine ratio*.

(2) is the definition of the *cosine ratio*.

(3) is the definition of the *tangent ratio*.

$$\text{By } T_4, \sin^2 + \cos^2 = 1.$$

$$\therefore a^2 = h^2 \sin^2, \text{ and } b^2 = h^2 \cos^2.$$

$$\therefore (4)_1 h^2 = a^2 + b^2; \quad (4)_2 \sin^2 + \cos^2 = 1 = \cot A \cot B.$$


(4)₁ is the *Pythagorean formula*.

$$(5) \text{ By } T_4, \sin A = \cos B, \quad \cos A = \sin B.$$

(5) is the *complementary formula*.

NOTE. For construction of figure, p. 13, see p. 15.

SPHERICAL.



THE FUNDAMENTAL ANALOGIES.

$$(1) \text{ By } T_1, \sin A = \frac{\sin a}{\sin h}, \sin a = \sin h \sin A, \sin h = \frac{\sin a}{\sin A}.$$

$$(2) \text{ By } T_3, \cos A = \frac{\tan b}{\tan h}, \tan b = \tan h \cos A, \tan h = \frac{\tan b}{\cos A}.$$

$$(3) \text{ By } T_2, \tan A = \frac{\tan a}{\sin b}, \tan a = \sin b \tan A, \sin b = \frac{\tan a}{\tan A}.$$

(1) is the sine analogy, or $\sin Ay$.

(2) is the cosine analogy, or $\cos Ay$.

(3) is the tangent analogy, or $\tan Ay$.

By T_4, VA_1H_1 , $\cos h = \cos a \cos b$, by (3).

$$(4) \cos h = \cos a \cos b = \cot A \cot B.$$

The first is the Pythagorean analogy.

Dividing $\sin A$ (1) by $\cos B$ (2), using (4).

$$(5) \sin A = \cos B : \cos b, \sin B = \cos A : \cos a.$$

(5) is the complementary analogy.

NOTE. If a sphere of unit radius be described about V as a centre, the faces will cut out a right spherical triangle having the sides a, b , and h .

NEGATIVE ANGLES.

NOTE. A point moving to the right generates a + distance, but moving back to the left tends to destroy this, and passing the origin generates a - distance.

A straight line revolving in the order of the quadrants I, II, III, IV, generates a + angle, but revolving back tends to destroy this, and passing the initial line generates a - angle.

RULE. *Changing the sign of an angle changes the sign of its sine, but not of its cosine.*

\therefore changes that of its tangent.

$+\angle$	I.	II.	III.	IV.	$-\angle$	-I.	-II.	-III.	-IV.
sin	+	+	-	-	sin	-	-	+	+
cos	+	-	-	+	cos	+	-	-	+
tan	+	-	+	-	tan	-	+	-	+

Since the terminus is changed across the initial line II' , but not across the transverse line TT' .


That is, $I' = IV$, $II' = III$, $III' = II$, $IV' = I$.

FUNCTIONS OF $nr \pm A$.

RULE. If an *acute* angle be added to or subtracted from an *even* number of quadrants, the functions of the resulting angle are equal arithmetically in value to the *like-named* functions of the acute angle; but if an acute angle be added to or subtracted from an *odd* number of quadrants, the functions of the resulting angle are arithmetically equal in value to the *co-named* functions of the acute angle.

By the equality of the *eight* possible ratio triangles, and the fact that for an even number of quadrants a and o are

the same as for A , but for an odd number they are interchanged.



NOTE. By revolving 1_A and 1_{-A} through any number of right angles, one rotation changes sine to cosine, two restores, and so on.

CONSTRUCTION.

(1) Lay off from the vertex V of a right trihedral a unit on each edge (VH being edge of rt. \angle).

(2) Through the extremity of one of the acute edges, as B_1 , pass a plane \perp to the other acute edge VA , thus:

Draw $BH \perp VH$, then $HA \perp VA$, lastly join AB .

(By Geom.) BAH is the *plane* measure of dihedral having edge VA .

(3) Through the other extremities H_2 and A_3 , pass planes \parallel to $A_1B_1H_1$, $\therefore \perp$ to VA .

(4) By p. 12, the parts of the nine right triangles are as given.

NOTE. Napier's Circular Parts are: The two sides about the right angle, the complements of the opposite angles, and the complement of the hypotenuse.

His rules are:

RULE I. The sine of the middle part is equal to the product of the tangents of the *adjacent* parts.

RULE II. The sine of the middle part is equal to the product of the cosines of the *opposite* parts.

$$\text{By } \begin{cases} \sin a = \sin A \sin h = \tan b \cot B \\ \sin b = \sin B \sin h = \tan a \cot A \end{cases} (3)$$

$$(5) \begin{cases} \cos A = \sin B \cos a = \tan b \cot h \\ \cos B = \sin A \cos b = \tan a \cot h \end{cases} (2)$$

$$(4) \cos h = \cos a \cos b = \cot A \cot B (4)$$

I. By (Comp. Ay.) An oblique angle and its opposite side are in the same quadrant.

II. By (P. Ay.) $h < 90^\circ$ when a and b are in the same quadrant.

$h > 90^\circ$ when a and b are in different quadrants.

EXAMPLES INVOLVING RIGHT TRIANGLES.

Solve in order of formulas. Each triangle furnishes nine examples.

	$s, c, ?$ h	$s, ?, t$ a	$?, c, t$ b	s, c, t A	$?, ?, ?$ B	
1.	10.40	5.43	8.87	$31^\circ 30'$	$58^\circ 30'$	$s = \sin \text{Def.}$
2.	37.57	27.81	25.26	$47^\circ 45'$	$42^\circ 15'$	$c = \cos \text{Def.}$
3.	7855.	6967.8	3627.4	$62^\circ 30'$	$27^\circ 30'$	$t = \tan \text{Def.}$
4.	50	30	40	$36^\circ 52' 10''$	$53^\circ 7' 30''$	$P = \text{Pyth. Theo.}$
5.	13	5	12	$22^\circ 37' 10''$	$67^\circ 22' 50''$	Comp.=co-relation.

6. The distance of the moon being h , and earth's radius a $A = 57' 2''$; find h . For the sun, $A = 8.8''$.

7. What is the length of the horizontal shadow of the Washington Monument, when the altitude of the sun is 50° ?

8. What is the radius of the circle of latitude on which you live?

9. (a) The angle of elevation of the top of a spire 500 feet distant is measured and found to be 13° . What is its height [above the instrument]?

(b) The elevation of base of spire is 9° . What is its height?

10. The elevation of the top of a spire is 45° , and at a point 100 feet farther away, $36^\circ 52'$. What is the height?

$$y = \frac{tt'}{t - t'}.$$

Spherical Right Triangles. First find a from A and h , then A from a and h . That is, solve in the order that the formulas are given (p. 13).

	a	b	h	A	B	
1.	$20^\circ 4' 22''$	$59^\circ 33' 43''$	$23^\circ 27' 29''$	S. Ay.
2.	$12^\circ 56' 43''$	$34^\circ 14' 45''$	$23^\circ 27' 31''$
3.	$57^\circ 21' 33''$	$59^\circ 33' 43''$	$23^\circ 27' 29''$	C. Ay.
4.	$43^\circ 20' 58''$	$44^\circ 9' 38''$	$13^\circ 34' 30''$
5.	$45^\circ 7' 49''$	$85^\circ 45' 2''$	$45^\circ 12' 33''$	T. Ay.
6.	$38^\circ 1' 40''$	$139^\circ 24'$	$50^\circ 14' 8''$
7.	$22^\circ 7' 15''$	$69^\circ 30' 13''$	$71^\circ 4' 20''$	P. Ay.
8.	$2^\circ 59'$	$22^\circ 21' 51''$	$22^\circ 33' 12''$
9.	$43^\circ 35'$	$46^\circ 59'$	$58^\circ 50'$	Comp. Ay.
10.	$22^\circ 15' 7''$	$27^\circ 28' 38''$	$73^\circ 27' 11''$

The right ascension R , declination d , and longitude L , of the sun form a right triangle of which these are b , a , and h ; A being the obliquity of the ecliptic.

11. $L = 214^\circ 14' 45''$; find R and d .

12. $R = 18$ hrs. 44 min. 50 sec.; find L and d .

13. $R = 4$ hrs. 38 min. 0.88 sec., $d = 22^\circ 7' 13.7''$; find L and A .

REMARKS AND QUESTIONS.

Many points rest directly upon page 12. Thus pages 13, 14, 18, 19, and 23 in great part.

As the last of 24, and most of 25, require 19 and 20, the given order has been followed.

It is very important for the student to observe as to what rests directly on pages 12 and 13.

Give the values of the functions of: 60° , 120° , 150° , 225° , -30° , -60° , -1210° , 350° , 440° , 1000° , 1234° .

How are the functions of 115° related to those of 205° ?

Reduce $(x^2 + y^2) \cos 720^\circ - 2xy \sin 540^\circ$.

When is it sufficient to consider angles in (1) I, (2) I and II, (3) I, -IV, (4) V, -VI,?

VALUES OF ONE FUNCTION IN TERMS OF ANOTHER.

By P. Theo., $1^2 = \sin^2 + \cos^2 = 1$. (1)

$\therefore \sin = \sqrt{1 - \cos^2}$, $\cos = \sqrt{1 - \sin^2}$.

NOTE. Prove this without P. Theo., and thus prove the theorem.

By P. Theo., $\left(\frac{1}{\cos}\right)^2 = \sec^2 = \tan^2 + 1$.

$\therefore \cos = \frac{1}{\sqrt{1 + \tan^2}}$, $\sin = \tan \cos = \frac{\tan}{\sqrt{1 + \tan^2}}$.

Give the value of each function in terms of the others.

1. $\sin 30^\circ = \frac{1}{2}$; find the other functions.

2. $\cos 45^\circ = \frac{1}{2}\sqrt{2}$; find the other functions.

3. $\tan A = 1, 2, 3, \dots$; find other functions.

HINT. If $\tan = 3$, $\sec \cos^2 = 10$. $\therefore \cos = \frac{1}{\sqrt{10}}$, $\sin = \frac{3}{\sqrt{10}}$.

4. Given one function of 90° ; find the others.

5. (1) $2 \sin = \cos$ (2) $\tan = 2 \sin$.

(3) $\sin = \cos$. (4) $\sin = \tan$.

HINT ON (2). $\frac{\sin}{\cos} = 2 \sin$. $\therefore \cos = \frac{1}{2}$. $\angle = 60^\circ$.


HINT ON (1). $\sin = x$, $2x = \sqrt{1 - x^2}$. $\therefore x = \sin = \frac{1}{5}\sqrt{5}$.

6. $\cos^2 \tan^2 + \sin^2 \cot^2 = 1$.

7. In rt. sph. \triangle , $\sin^2 h = \sin^2 o + \sin^2 a [\cos^2 o]$ (2d P. Ay.).
 $= \sin^2 a + \sin^2 o [\cos^2 a]$.

Its limit, $h^2 = a^2 + o^2$.

(a) FUNCTIONS OF SUM.



Directly from figure and page 12,

$$\sin(A+B) = \sin A \cos B + \cos A \sin B. \quad (1)$$

$$\cos(A+B) = \cos A \cos B - \sin A \sin B. \quad (2)$$

Dividing sine by cosine, then both terms of fractions by $\cos A \cos B$,

$$\tan(A+B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}. \quad (3)$$

(b) **Double Angle.** If $A = B$.

$$\sin 2A = 2 \sin A \cos A. \quad (1)$$

$$\cos 2A = \cos^2 A - \sin^2 A. \quad (2)$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}. \quad (3)$$

(c) **Sum and Difference of Sines.** If $A+B=S$, and $A-B=D$.

$$A = \frac{S}{2} + \frac{D}{2}, \quad B = \frac{S}{2} - \frac{D}{2}.$$

$$\therefore \sin A = \sin \frac{S}{2} \cos \frac{D}{2} + \cos \frac{S}{2} \sin \frac{D}{2}$$

$$\sin B = \sin \frac{S}{2} \cos \frac{D}{2} - \cos \frac{S}{2} \sin \frac{D}{2}$$

$$\sin A + \sin B = 2 \sin \frac{S}{2} \cos \frac{D}{2} \quad (1)$$

$$\sin A - \sin B = 2 \cos \frac{S}{2} \sin \frac{D}{2} \quad (2)$$

(a) FUNCTIONS OF DIFFERENCE.

$$\begin{aligned}\sin(A - B) &= \sin[A + (-B)] \\ &= \sin A \cos(-B) + \cos A \sin(-B).\end{aligned}$$

By rule for negative angles, page 14,

$$\sin(A - B) = \sin A \cos B - \cos A \sin B. \quad (1)$$

$$\cos(A - B) = \cos A \cos B + \sin A \sin B. \quad (2)$$

$$\tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}. \quad (3)$$

(b) **Half Angle.** By double angle formulas,

$$\cos^2 \frac{A}{2} + \sin^2 \frac{A}{2} = 1.$$

$$\cos^2 \frac{A}{2} - \sin^2 \frac{A}{2} = \cos A.$$

\therefore having the sum and the difference of \sin^2 and \cos^2 ,

$$\sin \frac{A}{2} = \sqrt{\frac{1 - \cos A}{2}}. \quad (1)$$

$$\cos \frac{A}{2} = \sqrt{\frac{1 + \cos A}{2}}. \quad (2)$$

“co” + sin -,

$$\tan \frac{A}{2} = \sqrt{\frac{1 - \cos A}{1 + \cos A}}. \quad (3)$$

(c) **Sum and Difference of Cosines.**

$$\cos A = \cos \frac{S}{2} \cos \frac{D}{2} - \sin \frac{S}{2} \sin \frac{D}{2}$$

$$\cos B = \cos \frac{S}{2} \cos \frac{D}{2} + \sin \frac{S}{2} \sin \frac{D}{2}$$

$$\cos A + \cos B = 2 \cos \frac{S}{2} \cos \frac{D}{2} \quad (1)$$


$$\cos A - \cos B = -2 \sin \frac{S}{2} \sin \frac{D}{2} \quad (2)$$

EXERCISES.

1. Find sine and cosine of $90^\circ \mp A$, $180^\circ \mp A$, ...
2. Find $\sin 3A = \sin(2A + A) = 3 \sin A - 4 \sin^3 A$,
 $\cos 3A = \cos(2A + A) = 4 \cos^3 A - 3 \cos A$.
- 3-7. If $\sin 30^\circ = \frac{1}{2}$, (3) find sine and cosine of 15° ;
 (4) of 45° ; (5) of $22\frac{1}{2}^\circ$; (6) of $67\frac{1}{2}^\circ$;
 (7) as answer, find those of 90° , or $(67\frac{1}{2}^\circ + 22\frac{1}{2}^\circ)$.
8. $\sin x + \cos x = \sqrt{1 + \sin 2x}$.
9. If $\angle \tan t$ be read "the angle whose tangent is t ," show that $\angle \tan \frac{1}{2} + \angle \tan \frac{1}{3} = 45^\circ$; also, $A + B = 90^\circ$ if $A = \angle \sin \frac{2}{3}$ and $B = \angle \sin \frac{4}{5}$.
10. Find the area of a regular dodecagon inscribed in a circle of radius 12.

ADDITIONAL.

1. If $a + b = \frac{1 + \sqrt{3}}{2}h$, find $\sin A$.
2. Find $\tan \frac{A}{2}$ by bisecting A .
 HINT. If the bisector divide $\sin A$ into two parts, x and y , $x : y = 1 : \cos A$.
 $\therefore \tan \frac{A}{2} = \frac{y}{\cos A} = \frac{x + y}{1 + \cos A} = \frac{\sin A}{1 + \cos A}$.
3. If $\tan \frac{a}{2} \tan \frac{b}{2} \tan \frac{c}{2} = 1$, find $\sin a$ and $\cos a$ in terms of the sine and cosine of b and c .
- 3'. If $\tan \frac{v}{2} \tan \left(45^\circ - \frac{P}{2}\right) \cot \frac{E}{2} = 1$, find the sine and cosine of each angle v , P , and E , in terms of those of the other two. (Due to Prof. O. Stone.)
- 3''. $\tan \left(45^\circ - \frac{P}{2}\right) = \sqrt{1 - \sin P} : \sqrt{1 + \sin P}$.



4. Find sum and difference of sines and cosines directly from figure and page 12.

HINT. The diagonals of a rhombus bisect at right angles. Half sum = greater — half difference.

$$\sin A + \sin B = 2 YY' = 2 \sin \frac{S}{2} VY'.$$

$$5. (\cos A - \cos B)^2 + (\sin A - \sin B)^2 = 4 \sin^2 \frac{A - B}{2}.$$

$$6. \cos 2A = (1 - \tan^2 A) : (1 + \tan^2 A).$$

$$7. \text{As } 2\pi r = 360^\circ, r^\circ = \frac{180^\circ}{\pi} = 57.3.$$

How many degrees in $1.5r$? *Ans.* $\frac{270^\circ}{\pi}$.

What is the length of 80° , r being the unit?

$$\text{Ans. } \frac{80}{57.3} = \frac{4}{9}\pi.$$

8. If $1_A = \cos A + \cdot \sin A$, $\therefore \cos A = \frac{1_A + 1_{-A}}{2}$, and $\sin A = \frac{1_A - 1_{-A}}{2}$, $1_A \times 1_B$ being $1_{(A+B)}$, and the binomial formula holding for such quantity, find $\sin(A+B)$, $\cos(A+B)$, $\sin A + \sin B$, etc., including $\cos 2A$, $\sin 3A$, $\cos 4A$, $+\cdot = +\sqrt{-1}$. $1_A =$ initial $\cos A +$ transverse $\sin A$.

9. The earth's radius subtends an angle of $57'$ at the moon; what is the distance (by 7)? The moon's apparent diameter is $31'$; what is it in miles (by 7)?

10. Construct the figure of page 13 :
- (1) When one side is in I, and the other in II.
 - (2) When both are in II, $\therefore h$ in I.
11. Construct the figure of page 19 :
- (1) When A and B are in I, but $(A + B)$ in II.
 - (2) When A is in I, and B in II.
 - (3) When both are in II.
12. Find $\tan (A + B)$ from the figure of page 19.
- HINT. The base of figure is now to be taken as 1.

NOTE. The "12 additional" are only for the leaders of the class, and not for the body.

Part of 8, though sometimes found in Algebra, seems *more nearly* in place here.

$$1_A = \cos A + \cdot \sin A, \quad (1)$$

$$1_B = \cos B + \cdot \sin B; \quad (2)$$

multiplying, $1_{(A+B)} = (\cos A \cos B - \sin A \sin B) +$
 $(\sin A \cos B + \cos A \sin B),$



but $1_{(A+B)} = \cos (A + B) + \cdot \sin (A + B).$

\therefore the usual formulas.

If instead of multiplying (1) and (2) we add them, observing (p. 22) that a journey of $2 \cos \frac{D}{2}$ in direction $\frac{S}{2}$ causes the same change of position as the two journeys 1_A and 1_B , we have the usual formulas for $\sin A + \sin B$ and $\cos A + \cos B$.

PLANE.

Law of Sines. In any plane triangle, the sides are proportional to the sines of the opposite angles.



By definition of sine,

$$a \sin B = p = b \sin A. \quad \therefore a : b = \sin A : \sin B.$$

Law of Cosines. The *square* of any side of a (pl.) triangle is equal to the *sum* of the *squares* of the other two sides, minus twice their product by the cosine of the included angle.

Pythagorean Proposition :

$$a^2 = p^2 + m^2, \quad b^2 = p^2 + n^2.$$

$$\therefore a^2 - b^2 = m^2 - n^2. \quad (\text{In words.})$$

$$a^2 - b^2 = (c - n)^2 - n^2 = c^2 - 2cn.$$

Definition of Cosine :

$$n = b \cos A.$$

\therefore substituting and transposing term b^2 ,

$$a^2 = b^2 + c^2 - 2bc \cos A.$$


Law of Tangents. The sum of any two sides of a (pl.) triangle is to their difference as the tangent of one-half the sum of the opposite angles is to the tangent of one-half their difference.

By law of sines and theory of proportion,

$$\frac{a+b}{a-b} = \frac{\sin A + \sin B}{\sin A - \sin B} = \frac{\tan \frac{1}{2}(A+B)}{\tan \frac{1}{2}(A-B)}.$$

SPHERICAL.

Law of Sines. In any spherical triangle, the sines of the sides are proportional to the sines of the opposite angles.



By sin Ay.,

$$\sin a \sin B = \sin p = \sin b \sin A.$$

$$\therefore \sin a : \sin b = \sin A : \sin B.$$

Law of Cosines. The cosine of any side of a (sph.) triangle is equal to the *product* of the *cosines* of the other two sides, plus the product of their sines by the cosine of their included angle.

Pythagorean Analogy:

$$\cos a = \cos p \cos m, \quad \cos b = \cos p \cos n.$$

$$\therefore \cos a : \cos b = \cos m : \cos n. \quad (\text{In words.})$$

$$\begin{aligned} \cos a : \cos b &= \cos(c-n) : \cos n \\ &= \cos c + \sin c \tan n. \end{aligned}$$

Cosine Analogy:

$$\tan n = \tan b \cos A.$$

Substituting and transposing factor $\cos b$,

$$\cos a = \cos b \cos c + \sin b \sin c \cos A.$$

For Analogy of Law of Tangents, see the limit of Napier's (2), page 29.

FUNCTIONS OF HALF ANGLES.

Plane. By law of cosines,

$$\cos A = \frac{a^2 - (b^2 + c^2)}{-2bc} = \frac{b^2 + c^2 - a^2}{2bc}.$$

PROBLEM. To substitute this value in $1 \pm \cos A$, and thus find the functions of the half angles in terms of the sides.

If $a + b + c = 2s$,

$$\therefore a + b - c = 2(s - c),$$

$$a - b + c = 2(s - b),$$

$$b + c - a = 2(s - a).$$

$$\begin{aligned} 1 - \cos A &= \frac{a^2 - b^2 - c^2 + 2bc}{2bc} = \frac{a^2 - (b - c)^2}{2bc} \\ &= \frac{2(s - b) 2(s - c)}{2bc}. \end{aligned}$$

$$\text{Similarly, } 1 + \cos A = \frac{2s(s - a)}{bc}.$$

$$\therefore \sin \frac{A}{2} = \sqrt{\frac{(s - b)(s - c)}{bc}}; \text{ from } \sin \frac{A}{2} = \sqrt{\frac{1 - \cos A}{2}}.$$

$$\cos \frac{A}{2} = \sqrt{\frac{s(s - a)}{bc}}; \quad \text{from } \cos \frac{A}{2} = \sqrt{\frac{1 + \cos A}{2}}.$$

$$\tan \frac{A}{2} = \sqrt{\frac{(s - b)(s - c)}{s(s - a)}}.$$

That is, the sine of half an angle of a plane triangle is equal to the square root of half the sum of the three sides minus one of the including sides, into the half sum minus the other including side, divided by the product of the including sides.

Give for cosine and tangent.

Ex. Find the value of $\tan \frac{A}{2}$ directly from page 12, and Wentworth's Geometry, page 250.

If r be the radius of inscribed circle, $\text{area} = rs$.

By Geometry,

$$\text{area} = \sqrt{s(s-a)(s-b)(s-c)}.$$

$$\therefore r = \sqrt{\frac{(s-a)(s-b)(s-c)}{s}}.$$

By figure, $\tan \frac{A}{2} = \frac{r}{s-a}.$

FORMULAS FOR HALF ANGLES.

Spherical. By law of cosines,

$$\cos A = \frac{\cos a - \cos b \cos c}{\sin b \sin c}.$$

As in Plane,

$$\begin{aligned} 1 - \cos A &= \frac{\sin b \sin c + \cos b \cos c - \cos a}{\sin b \sin c} \\ &= \frac{\cos a - \cos(b-c)}{-\sin b \sin c} \quad (\text{dif. of cos}) \\ &= \frac{2 \sin(s-b) \sin(s-c)}{\sin b \sin c}. \end{aligned}$$

Similarly,

$$1 + \cos A = \frac{2 \sin s \sin(s-a)}{\sin b \sin c}.$$

\therefore the analogy here is the same as in law of sines,

$$\sin \frac{A}{2} = \sqrt{\frac{\sin(s-b) \sin(s-c)}{\sin b \sin c}},$$

$$\cos \frac{A}{2} = \sqrt{\frac{\sin s \sin(s-a)}{\sin b \sin c}},$$

$$\tan \frac{A}{2} = \sqrt{\frac{\sin(s-b) \sin(s-c)}{\sin s \sin(s-a)}}.$$

That is, the cosine of one-half of either angle of a spherical triangle is equal to the square root of the sine of one-half of the sum of the three sides, into the sine of one-half this

sum minus the side opposite the angle, divided by the product of the sines of the including sides.

Give sine and tangent in words.

LAW OF COSINES FOR ANGLES. FUNCTIONS OF HALF SIDES.

If $A'B'C'$ be the polar of ABC , by first law of cosines and Geometry,

$$\cos A = -\cos B \cos C + \sin B \sin C \cos a.$$

In words,


$$\cos a = \frac{\cos A + \cos B \cos C}{\sin B \sin C},$$

$$\sin \frac{a}{2} = \sqrt{\frac{-\cos S \cos (S - A)}{\sin B \sin C}},$$

$$\cos \frac{a}{2} = \sqrt{\frac{\cos (S - B) \cos (S - C)}{\sin B \sin C}}.$$

In words. Note Analogies.

THE GAUSS EQUATIONS.



$$\sin \frac{1}{2}(A + B) \cos \frac{1}{2}c = \cos \frac{1}{2}(a - b) \cos \frac{1}{2}C.$$

$$\cos \frac{1}{2}(A + B) \cos \frac{1}{2}c = \cos \frac{1}{2}(a + b) \sin \frac{1}{2}C.$$

$$\sin \frac{1}{2}(A - B) \sin \frac{1}{2}c = \sin \frac{1}{2}(a - b) \cos \frac{1}{2}C.$$

$$\cos \frac{1}{2}(A - B) \sin \frac{1}{2}c = \sin \frac{1}{2}(a + b) \sin \frac{1}{2}C.$$

$$f \frac{1}{2}(A \pm B) \quad f \frac{1}{2}c = f \frac{1}{2}(a \pm b) \quad f \frac{1}{2}C.$$

RULE I. \sin in (1) gives $-$ in (3) and conversely.
 \cos in (1) gives $+$ in (3) and conversely.

RULE II. Functions have same names in (2) and (3).
 Functions have co-names in (4) and (1).

NOTE. The rules also hold for I, II, III, and IV.

To prove (1), $\sin \frac{1}{2}(A + B) = \sin \left(\frac{A}{2} + \frac{B}{2} \right)$.

$$\begin{aligned}
 \text{Now } \sin \left(\frac{A}{2} + \frac{B}{2} \right) &= \sin \frac{A}{2} \cos \frac{B}{2} + \cos \frac{A}{2} \sin \frac{B}{2} \\
 &= \sqrt{\frac{\sin(s-b) \sin(s-c)}{\sin b \sin c}} \sqrt{\frac{\sin s \sin(s-b)}{\sin a \sin c}} \\
 &\quad + \sqrt{\frac{\sin s \sin(s-a)}{\sin b \sin c}} \sqrt{\frac{\sin(s-a) \sin(s-c)}{\sin a \sin c}} \\
 &= \frac{\sin(s-b)}{\sin c} \cos \frac{C}{2} + \frac{\sin(s-a)}{\sin c} \cos \frac{C}{2} \\
 &= \frac{\sin(s-b) + \sin(s-a)}{\sin c} \cos \frac{C}{2} \quad (\text{by sum of sines}) \\
 &= \frac{2 \sin \left(s - \frac{a+b}{2} \right) \cos \frac{a-b}{2}}{\sin c} \cos \frac{C}{2};
 \end{aligned}$$

$$\text{but } s - \frac{a+b}{2} = \frac{c}{2},$$

$$\therefore = \frac{2 \sin \frac{c}{2} \cos \frac{a-b}{2}}{2 \sin \frac{c}{2} \cos \frac{c}{2}} \cos \frac{C}{2}. \quad \therefore \text{Q. E. D.}$$

By the Gauss Equations and (Div. Ax.) we have

NAPIER'S PROPORTIONS.

- (1) $\sin \frac{1}{2}(a + b) : \sin \frac{1}{2}(a - b) :: \cot \frac{1}{2}C : \tan \frac{1}{2}(A - B)$.
- (2) $\cos \frac{1}{2}(a + b) : \cos \frac{1}{2}(a - b) :: \cot \frac{1}{2}C : \tan \frac{1}{2}(A + B)$.
- (3) $\sin \frac{1}{2}(A + B) : \sin \frac{1}{2}(A - B) :: \tan \frac{1}{2}c : \tan \frac{1}{2}(a - b)$.
- (4) $\cos \frac{1}{2}(A + B) : \cos \frac{1}{2}(A - B) :: \tan \frac{1}{2}c : \tan \frac{1}{2}(a + b)$.

THEOREM 1. *The sine of one-half the sum of either two sides of any spherical triangle is to the sine of one-half their difference as the cotangent of one-half the angle which they include is to the tangent of one-half the difference of the angles opposite.*

THEOREM 2. *The cosine of one-half the sum of either two sides of any spherical triangle is to the cosine of one-half their difference as the cotangent of one-half the angle which they include is to the tangent of one-half the sum of the angles opposite.*

THEOREM 3. *The sine of one-half the sum of either two angles of any spherical triangle is to the sine of one-half their difference as the tangent of one-half the side which they include is to the tangent of one-half the difference of the sides opposite.*

THEOREM 4. *The cosine of one-half the sum of either two angles of any spherical triangle is to the cosine of one-half their difference as the tangent of one-half the side which they include is to the tangent of one-half the sum of the sides opposite.*

Note the analogies: sine giving —, cosine, +; side giving “co” *f*, angle giving function.

EXAMPLES UNDER OBLIQUE TRIANGLES.

Given (any three parts, one being a side) :

- I. One side and two angles, L. of Ss.
 - II. Two sides and \angle opposite one of them, L. of Ss.
 - III. Two sides and the included \angle , L. of Ts. or L. of Cs.
 - IV. Three sides. Formulas for half \angle s or L. of Cs.
 - V. Sph. Three \angle s. Formulas for half sides.
- I. $(a, A, B) (b, A, B) (c, A, B) (a, A, C) (b, A, C)$
 $(c, A, C) (a, B, C) (b, B, C) (c, B, C).$
 - II. $(a, b, A) (a, b, B) (a, c, A) (a, c, C) (b, c, B) (b, c, C).$
 - III. $(a, b, C) (a, c, B) (b, c, A).$

	<i>a</i>	<i>b</i>	<i>c</i>	<i>A</i>	<i>B</i>	<i>C</i>
3.	1686	960	2400	33° 34' 39"	18° 21' 21"	128° 4'
4.	40	34	45	83° 53' 15"	57° 41' 25"	38° 25' 20"
5.	10	12	14	44° 24' 55"	57° 7' 18"	78° 27' 47"
1.	6	8	10	36° 52'	53° 8'	90°
2.	2	$\sqrt{6}$		45°	60°	75°

6. Find the distance between two objects (*supposed inaccessible*) by calculating the distance to each by right triangles, also by law of sines, then measuring the angle between these distances. Measure the distance to test your answer.

7. If in (4) the unit be one mile, and CA be east and west line, what is the direction of each vertex from the others?

8. An object when viewed from the ends of an east and west line, of length 34, bears $N. 51^{\circ} 35' W.$ and $N. 6^{\circ} 7' E.$ What is its distance from each, and from the straight line joining them?

	a	b	c	A	B	C
1.	$70^{\circ} 4' 17''$	$63^{\circ} 21' 26''$	$58^{\circ} 16' 22''$	$81^{\circ} 38' 20''$	$70^{\circ} 9' 38''$	$63^{\circ} 31' 20''$
2.	$20^{\circ} 16'$	$56^{\circ} 18'$	$66^{\circ} 18'$	$20^{\circ} 10'$	$55^{\circ} 54'$	$114^{\circ} 18' 30''$
3.	$147^{\circ} 5' 33''$	$155^{\circ} 5' 18''$	$33^{\circ} 1' 36''$	$110^{\circ} 10'$	$133^{\circ} 18'$	$70^{\circ} 20' 40''$
4.	$69^{\circ} 34' 56''$		$70^{\circ} 20' 20''$	$50^{\circ} 10' 10''$		$50^{\circ} 30' 8''$

As the chief applications are to Astronomy, we give

THE FIRST ASTRONOMICAL TRIANGLE PZH .

P is north pole of equator.

Z is zenith (pole of horizon).

H is the heavenly body.


Altitude of pole = NP = latitude of place.

The distances of a body from any great circle and its pole are complementary.

PZ = co-latitude. HP = polar distance = co-declination.

HZ = zenith distance = co-altitude.

$\angle Z$ = azimuth. $\angle P$ = hour or time angle.




EXAMPLE. Calculate the following for a place, latitude 45° , and for the longest day of the year.

- (a) The azimuth of sun at setting and rising.
 (b) Time of rising and setting.
 (c) The greatest altitude of the sun.
 (d) Time when a vertical object casts a horizontal shadow of its own length.

THE SECOND ASTRONOMICAL TRIANGLE.

Before considering this triangle involving two great circles and their poles, we will consider a point referred to *any* great circle and its pole.

Let V (any point of this great circle) be the origin, and consider the hemisphere limited by the great circle of V .



Let P be the pole, and EQ the great circle of reference, H the heavenly body; VH is, by Geometry, \perp to great circle of V cutting it in G ; denote PG by b , HG by a .

VF = direct co-ordinate D , FQ = co. D . (Fig. 2.)

FH = transverse co-ordinate T , HP = co. T .

From the rt. $\triangle HPG$ and HVF (C. Ay.),

$$\tan b = \cot T \sin D. \quad (1)$$

$$\therefore \tan T = \cot b \sin D. \quad (2)$$

$$(\text{Tan. Ay.}), \quad \tan(\text{co. } D) = \cot D = \frac{\tan a}{\sin b}. \quad (3)$$

$$(\text{P. Ay.}), \quad \cos D \cos T = \cos VH. \quad (4)$$

In the second triangle $PP'H$, we have two systems of co-ordinates: Right ascension (R) and declination (d), when

referred to the equator EQ , but latitude l and longitude L , when referred to the ecliptic EC . Since, if the former be D and T , the latter are D' and T' , if V , the common point of the two circles, be the origin :

$$\tan b = \cot d \sin R. \quad (1)$$

$$\frac{\sin b}{\sin b'} = \frac{\tan R}{\tan L}. \quad (2) \text{ by } (3) \text{ above.}$$

$$\tan b' = \cot l \sin L. \quad (3)$$

$$\cos l \cos L = \cos d \cos R. \quad (4)$$

$PP' = E$, the obliquity of the ecliptic ; $P' = \cos L$; $P = 90 + R$.

The reduction of an observation from the equator to the ecliptic is of so much importance that one example is worked through and back.

<i>R</i>	81° 48' 42.4''	<i>L</i>	85° 45' 2.00''
<i>d</i>	68° 27' 54.5''	<i>l</i>	45° 7' 48.98''
sin <i>R</i>	9.9955499	sin <i>L</i>	9.9988044
cot <i>d</i>	9.5961411	cot <i>l</i>	9.9980251
tan <i>b'</i>	9.5916910	tan <i>b</i>	9.9968295
<i>b'</i>	21° 20' 1.58''	<i>b</i>	44° 47' 27.11''
<i>e</i>	23° 27' 25.53''	<i>e</i>	23° 27' 25.53''
<i>b</i> = <i>b'</i> + <i>e</i>	44° 47' 27.11''	<i>b'</i> = <i>b</i> - <i>e</i>	21° 20' 1.58''
sin <i>b</i>	9.8478940	sin <i>b'</i>	9.5608631
1 : sin <i>b'</i>	0.4391369	1 : sin <i>b</i>	0.1521060
tan <i>R</i>	0.8419622	tan <i>L</i>	1.1289931
tan <i>L</i>	1.1289931	tan <i>R</i>	0.8419622
<i>L</i>	85° 45' 2.00''	<i>R</i>	81° 48' 42.4''
sin <i>L</i>	9.9988044	sin <i>R</i>	9.9955499
cot <i>b</i>	0.0031705	cot <i>b'</i>	0.4083090
tan <i>l</i>	0.0019749	tan <i>d</i>	0.4038589
<i>l</i>	45° 7' 48.98''	<i>d</i>	68° 27' 54.5''
cos <i>l</i>	9.8484953	cos <i>d</i>	9.5647190
cos <i>L</i>	9.8698113	cos <i>R</i>	9.1535877
cos <i>l</i> cos <i>L</i>	8.7183066	cos <i>d</i> cos <i>R</i>	8.7183067

QUADRANTAL TRIANGLES.

By Geometry, the polar of a right triangle is a quadrantal triangle. If $A'B'C'$ be a rt. \triangle , and ABC its polar, then, by Napier's Rules,

$$\sin A = \tan B \cot b = \sin a \sin H.$$

$$\sin B = \tan A \cot a = \sin b \sin H.$$

$$\cos a = -\tan B \cot H = \cos A \sin b.$$

$$\cos b = -\tan A \cot H = \cos B \sin a.$$

$$\cos H = -\cot a \cot b = -\cos A \cos B.$$

That is, Napier's Rules would hold for a quadrantal triangle, if the circular parts be the two angles about the quadrant, the complements of the opposite sides, and the complement of the hypotenuse angle, giving

THREE SECONDARY ANALOGIES.

$$\sin a = \frac{\sin A}{\sin H}, \quad \cos a = -\frac{\tan B}{\tan H}, \quad \tan a = \frac{\tan A}{\sin B}.$$

RULE. Those formulas that contain a co-function of H have a negative sign.

I. The same as for right triangles.

Proof by I, of rt. \triangle , if $A \leq 90^\circ$, $a' \geq 90^\circ$, $A' \geq 90^\circ$, $\therefore a \leq 90^\circ$.

II'. The reverse of II, for right triangles.

PROOF. If A and B in same quadrant, $\therefore a'$ and b' in same.

By II, $h' < 90^\circ$, $\therefore H > 90^\circ$.

Q. E. D.

PROOF OF RULE. There are but two hypotheses :

(a) A and B in the same quadrant ;

(b) A and B in different quadrants.

If (a), by II'', $H > 90^\circ$, \therefore co-function $H -$.

But, by I', the other two factors are alike in signs.

\therefore sign of formula must be $-$.

If (b), by II, $H < 90^\circ$, \therefore co-function of H +.

But by I, the other two factors are alike in sign.

\therefore sign of formula must be —. \therefore Q. E. D.

EXAMPLES UNDER QUADRANTAL TRIANGLES.

1. In what latitude does the sun rise in the northeast at summer solstice?


2. Find time of sunrise and sunset at your school on the longest day of the year.

3. Rt. \triangle . Form a table of times when the shadow of the side of the east school door-way will coincide with an east and west crack in the hall floor.

CLOSING NOTE. Establish the definitions of sine and cosine without similar triangles, thence laws of sine and cosine and Pythagorean proposition. Then give trigonometric proofs of proportions concerning oblique lines, sides of a triangle opposite equal angles, and converse.

Greater side opposite greater angle, and converse.

Three propositions concerning similar triangles, and three concerning equal triangles, also any other trigonometric proofs of common geometric principles, thus tying the two subjects together.



MODERN LANGUAGES, Etc

MODERN LANGUAGES.

		INTROD.	PRICE
Becker :	Spanish Idioms		\$ 1.80
Collar :	Eysenbach's German Lessons		1.20
Cook :	Table of German Prefixes and Suffixes05
Doriot :	Illustrated Beginners' Book in French80
Knapp :	Modern French Readings80
	Modern Spanish Readings		1.50
	Modern Spanish Grammar		1.50
Spiers :	French-English General Dictionary		4.50
	English-French General Dictionary		4.50
Stein :	German Exercises40

MISCELLANEOUS.

Ariel :	Those Dreadful Mouse Boys75
Arrowsmith :	Kaegi's Rigveda		1.50
Burgess :	The American University15
Channing :	Delbrück's Introduction to the Study of Language94
Culver :	Epitome of Anatomy20
Davidson :	The Place of Art in Education20
	The Parthenon Frieze and other Essays		1.50
	A Dante Handbook		1.12
Dippold :	Emanuel Geibel's Brunhild45
Fisk :	Teachers' Improved Class Books		
Flagg :	Pedantic Versicles75
Halsey :	Genealogical and Chronological Chart25
Harrington :	Helps to the Intelligent Study of College Prepara- tory Latin		
Hitchcock :	Manual for the Gymnasium25
Hofmann :	Inaugural Address25
Hudson :	Daniel Webster25
J. B. G. :	Queen of Hearts20
Lectures on	School Hygiene80
Leighton :	Harvard Examination Papers		1.00
Prince :	Courses of Studies and Methods of Teaching75
Monoyer :	Sight Test12
Packard :	Studies in Greek Thought80
Seelye :	Growth through Obedience25
Smith & Blackwell :	Parallel Syntax Chart		1.00
Stevens :	Yale Examination Papers75
Straight :	True Aim of Industrial Education10
Super :	Weil's Order of Words in the Ancient Languages compared with the Modern		1.12
Warren :	True Key to Ancient Cosmology20

*Copies sent to Teachers for Examination, with a view to Introduction,
on receipt of Introduction Price.*

GINN & COMPANY, Publishers.

BOSTON.

NEW YORK.

CHICAGO.

SCIENCE AND HISTORY.

NATURAL SCIENCE.

		INTROD.	PRICE
Everett:	Vibratory Motion and Sound		\$ 2.00
Gage:	Elements of Physics		1.12
	Introduction to Physical Science.		1.00
Hale:	Little Flower-People40
Hill:	Questions on Stewart's Physics35
Journal of	Morphology	(per vol.)	6.00
Knight:	Primer of Botany30
Williams:	Introduction to Chemical Science80

PHILOSOPHICAL SCIENCE.

Davidson:	Rosmini's Philosophical System		2.50
Hickok:	Philosophical Works00
Ladd:	Lotze's Outlines of Metaphysic80
	Lotze's Outlines of Philosophy of Religion80
	Lotze's Outlines of Practical Philosophy80
	Lotze's Outlines of Psychology80
	Lotze's Outlines of Aesthetics80
	Lotze's Outlines of Logic80
Seelye:	Hickok's Mental Science (Empirical Psychology)		1.12
	Hickok's Moral Science		1.12

POLITICAL SCIENCE.

Clark:	Philosophy of Wealth		1.00
Clark & Giddings:	The Modern Distributive Process	(retail)	.75
Macy:	Our Government70
Political Science	Quarterly	(per vol.)	3.00
Seligman:	Railway Tariffs and the Interstate Law	(retail)	.75

HISTORY.

Allen:	Readers' Guide to English History25
Andrade:	Historia do Brazil75
Fiske-Irving:	Washington and His Country		1.00
Halsey:	Genealogical and Chronological Chart25
Journal of Archæology	(per vol.)	5.00
Judson:	Cæsar's Army		1.00
Montgomery:	Leading Facts of English History		1.00
	English History Reader60
Moore:	Pilgrims and Puritans60
Myers:	Mediæval and Modern History		1.50
	Ancient History		1.40

*Copies sent to Teachers for Examination, with a view to Introduction,
on receipt of Introduction Price.*

GINN & COMPANY, Publishers.

BOSTON.

NEW YORK.

CHICAGO.

Latin Text-Books.

	INTROD. PRICE.
ALLEN & GREENOUGH: Latin Grammar	\$1.12
Cæsar (7 books, with vocabulary; illustrated)	1.25
Cicero (13 orations, with vocabulary; illustrated)	1.25
Sallust's Catiline60
Cicero de Senectute50
Ovid (with vocabulary)	1.40
Preparatory Course of Latin Prose	1.40
Latin Composition	1.12
ALLEN New Latin Method90
Introduction to Latin Composition90
Latin Primer90
Latin Lexicon90
Remnants of Early Latin75
Germania and Agricola of Tacitus	1.00
BLACKBURN . . Essentials of Latin Grammar70
Latin Exercises60
Latin Grammar and Exercises (in one volume)	1.00
COLLAR & DANIELL: Beginner's Latin Book	1.00
Latine Reddenda (paper)20
Latine Reddenda and Voc. (cloth)30
COLLEGE SERIES OF LATIN AUTHORS.	
Greenough's Satires and Epistles of Horace (text edition) \$0.20; (text and notes)	1.25
CROW Selections from the Latin Poets	1.40
CROWELL & RICHARDSON: Brief History of Roman Lit. (BENDER)	1.00
GREENOUGH . . Virgil:—	
Bucolics and 6 Books of Æneid (with vocab.)	1.60
Bucolics and 6 Books of Æneid (without vocab.)	1.12
Last 6 Books of Æneid, and Georgics (with notes)	1.12
Bucolics, Æneid, and Georgics (complete, with notes)	1.60
Text of Virgil (complete)75
Vocabulary to the whole of Virgil	1.00
GINN & Co. . . Classical Atlas and Geography (cloth)	2.00
HALSEY. . . . Etymology of Latin and Greek	1.12
KEEP Essential Uses of the Moods in Greek and Latin25
KING Latin Pronunciation25
LEIGHTON. . . Latin Lessons	1.12
First Steps in Latin	1.12
MADVIG . . . Latin Grammar (by THACHER)	2.25
PARKER & PREBLE: Handbook of Latin Writing50
PREBLE. . . . Terence's Adelphoe25
SHUMWAY. . . Latin Synonyms30
STICKNEY. . . Cicero de Natura Deorum	1.40
TETLOW . . . Inductive Latin Lessons	1.12
TOMLINSON . . Manual for the Study of Latin Grammar20
Latin for Sight Reading	1.00
WHITE (J. W.) Schmidt's Rhythmic and Metric	2.50
WHITE (J. T.) Junior Students' Latin-English Lexicon (mor.)	1.75
English-Latin Lexicon (sheep)	1.50
Latin-English and English-Latin Lexicon (sheep)	3.00
WHITON . . . Auxilia Vergiliana; or, First Steps in Latin Prosody15
Six Weeks' Preparation for Reading Cæsar40

*Copies sent to Teachers for Examination, with a view to Introduction,
on receipt of Introduction Price.*

GINN & COMPANY, Publishers,
BOSTON, NEW YORK, AND CHICAGO.

GREEK BOOKS.²

Allen	Medea of Euripides	\$1.00
College Series of Greek Authors.... See D'Ooge, Dyer, Humphreys.		
D'Ooge	Sophocles' Antigone: <i>Text and Notes</i>95
	<i>Text only</i>45
Dyer	Plato's Apology and Crito: <i>Text and Notes</i>95
	<i>Text only</i>45
Flagg.....	Hellenic Orations of Demosthenes	1.00
	Anacreontics35
	Seven against Thebes	1.00
Goodwin	Greek Grammar	1.50
	Greek Reader	1.50
	Greek Moods and Tenses	1.50
	Selections from Xenophon and Herodotus	1.50
Goodwin & White:	Anabasis	1.00
	Anabasis (<i>with Vocabulary</i>)	1.50
Humphreys	Aristophanes' Clouds: <i>Text and Notes</i>95
	<i>Text only</i>45
Keep	Essential Uses of the Moods25
Kendrick	Greek at Sight15
Leighton	New Greek Lessons	1.20
Liddell & Scott..	Abridged Greek-English Lexicon	1.90
	Unabridged Greek-English Lexicon	9.40
Seymour	Selected Odes of Pindar	1.40
Sidgwick	Greek Prose Composition	1.50
Tarbell .	Philippics of Demosthenes	1.00
Tyler	Selections from Greek Lyric Poets	1.00
White	First Lessons in Greek	1.20
	Schmidt's Rhythmic and Metric of the Classical Languages	2.50
	Œdipus Tyrannus of Sophocles	1.12
	Stein's Dialect of Herodotus10
Whiton	Orations of Lysias	1.00

Copies sent to teachers for examination, with a view to Introduction, on receipt of Introduction Price given above.

GINN & COMPANY, Publishers.

BOSTON.

NEW YORK.

CHICAGO.

BOOKS ON ENGLISH LITERATURE.

Allen . . .	Reader's Guide to English History	\$.25
Arnold . . .	English Literature	1.50
Bancroft . .	A Method of English Composition50
Browne . . .	Shakespeare Versification25
Fulton & Trueblood :	Choice Readings	1.50
	Chart Illustrating Principles of Vocal Expression,	2.00
Genung . . .	Practical Elements of Rhetoric	1.25
Gilmore . . .	Outlines of the Art of Expression60
Ginn	Scott's Lady of the Lake	<i>Bds.</i> , .35; <i>Cloth</i> , .50
	Scott's Tales of a Grandfather	<i>Bds.</i> , .40; <i>Cloth</i> , .50
Gummere . .	Handbook of Poetics	1.00
Hudson . . .	Harvard Edition of Shakespeare: —	
	20 Vol. Edition. <i>Cloth, retail</i>25.00
	10 Vol. Edition. <i>Cloth, retail</i>	20.00
	Life, Art, and Character of Shakespeare. 2 vols.	
	<i>Cloth, retail</i>	4.00
	New School Shakespeare. <i>Cloth</i> . Each Play45
	Old School Shakespeare, per play20
	Expurgated Family Shakespeare	10.00
	Essays on Education, English Studies, etc.25
	Three Volume Shakespeare, per vol.	1.25
	Text-Book of Poetry	1.25
	Text-Book of Prose	1.25
	Pamphlet Selections, Prose and Poetry15
	Classical English Reader	1.00
Johnson . .	Rasselas	<i>Bds.</i> , .30; <i>Cloth</i> , .40
Lee	Graphic Chart of English Literature25
Martineau . .	The Peasant and the Prince	<i>Bds.</i> , .35; <i>Cloth</i> , .50
Minto	Manual of English Prose Literature	1.50
	Characteristics of English Poets	2.00
Rolfe	Craik's English of Shakespeare90
Scott	Guy Mannering	<i>Bds.</i> , .60; <i>Cloth</i> , .75
	Ivanhoe	<i>Bds.</i> , .60; <i>Cloth</i> , .75
	Talisman	<i>Bds.</i> , .50; <i>Cloth</i> , .60
	Rob Roy	<i>Bds.</i> , .60; <i>Cloth</i> , .75
Sprague . . .	Milton's Paradise Lost, and Lycidas45
	Six Selections from Irving's Sketch-Book	<i>Bds.</i> , .25; <i>Cloth</i> , .35
Swift	Gulliver's Travels	<i>Bds.</i> , .30; <i>Cloth</i> , .40
Thom	Shakespeare and Chaucer Examinations00

Copies sent to Teachers for Examination, with a view to Introduction,
on receipt of the Introduction Price given above.

GINN & COMPANY, Publishers,
Boston, New York, and Chicago.

Mathematics.²

*Introd.
Prices.*

Byerly.	Differential Calculus.....	\$2.00
	Integral Calculus.....	2.00
Ginn.	Addition Manual15
Halsted.	Mensuration	1.00
Hardy	Quaternions	2.00
Hill	Geometry for Beginners	1.00
Sprague	Rapid Addition10
Taylor	Elements of the Calculus	1.80
Wentworth	Grammar School Arithmetic75
	Shorter Course in Algebra.....	1.00
	Elements of Algebra	1.12
	Complete Algebra	1.40
	Plane Geometry75
	Plane and Solid Geometry	1.25
	Plane and Solid Geometry, and Trigonometry	1.40
	Plane Trigonometry and Tables. <i>Paper.</i> ..	.60
	Pl. and Sph. Trig., Surv., and Navigation .	1.12
	Pl. and Sph. Trig., Surv., and Tables.....	1.25
	Trigonometric Formulas	1.00
Wentworth & Hill:	Practical Arithmetic	1.00
	Abridged Practical Arithmetic75
	Exercises in Arithmetic	
	Part I. <i>Exercise Manual</i>	
	Part II. <i>Examination Manual</i>35
	Answers (to both Parts)25
	Exercises in Algebra70
	Part I. <i>Exercise Manual</i>35
	Part II. <i>Examination Manual</i>35
	Answers (to both Parts)25
	Exercises in Geometry70
	Five-place Log. and Trig. Tables (<i>7 Tables</i>)	.50
	Five-place Log. and Trig. Tables (<i>Comp. Ed.</i>)	1.00
Wentworth & Reed:	First Steps in Number, <i>Pupils' Edition</i>	.30
	<i>Teachers' Edition</i> , complete	.90
	Parts I., II., and III. (separate), each	.30
Wheeler	Plane and Spherical Trig. and Tables.....	1.00

*Copies sent to Teachers for examination, with a view to Introduction.
on receipt of Introduction Price.*

GINN & COMPANY, Publishers.

BOSTON.

NEW YORK.

CHICAGO.

Wentworth's Trigonometries.

THE aim has been to furnish just so much of Trigonometry as is actually taught in our best schools and colleges. The principles have been unfolded with the utmost brevity consistent with simplicity and clearness, and interesting problems have been selected with a view to awaken a real love for the study. Much time and labor have been spent in devising the simplest proofs for the propositions, and in exhibiting the best methods of arranging the logarithmic work. Answers are included.

Plane and Solid Geometry, and Plane Trigonometry.

12mo. Half morocco. 490 pages. Mailing Price, \$1.55; Introduction, \$1.40; Allowance for old book, 40 cents.

Plane Trigonometry.

12mo. Paper. 80 pages. Mailing Price, 35 cents; Introduction, 30 cents.

Plane Trigonometry Formulas.

Two charts (30 × 40 inches each) for hanging on the walls of the classroom. Introduction Price, \$1.00 per set.

Plane Trigonometry and Logarithms.

8vo. Paper. 160 pages. Mailing Price, 65 cents; Introduction, 60 cents.

Plane and Spherical Trigonometry.

12mo. Half morocco. iv + 151 pages. Mailing Price, 80 cents; for introduction, 75 cents.

Wentworth's Plane and Spherical Trigonometry, and Surveying.

With Tables. 8vo. Half morocco. 307 pages. Mailing Price, \$1.40; Introduction, \$1.25; Allowance for old book, 40 cents.

Surveying.

8vo. 80 pages. Paper. Mailing Price, 35 cents; for introduction, 30 cents.

Wentworth's Plane and Spherical Trigonometry, Surveying, and Navigation.

Surveying, and Navigation.

12mo. Half morocco. 330 pages. Mailing Price, \$1.25; Introduction, \$1.12; Allowance for old book, 40 cents.

THE object of the work on Surveying and Navigation is to present these subjects in a clear and intelligible way, according to the best methods in actual use; and also to present them in so small a compass, that students in general may find the time to acquire a competent knowledge of these very interesting and important studies. Answers are included.

S. J. Kirkwood, *Prof. of Mathematics, University of Wooster, O.*: Wentworth's Algebra, Geometry, and Trigonometry are excellent textbooks. (Dec. 15, 1883.)

Otis H. Robinson, *Prof. of Mathematics, University of Rochester*: I think Wentworth's Surveying an admirable introduction to the study of the subject. (May 28, 1883.)

Wentworth & Hill's Five-Place Logarithmic and Trigonometric Tables.

By G. A. WENTWORTH, A.M., and G. A. HILL, A.M.

Seven Tables (for Trigonometry and Surveying): Cloth. 8vo. 79 pages. Mailing Price, 55 cents; Introduction, 50 cents.

Complete (for Trigonometry, Surveying, and Navigation): Half morocco. 8vo. 158 pages. Mailing Price, \$1.10; Introduction, \$1.00.

THESE Tables have been prepared mainly from Gauss's Tables, and are designed for the use of schools and colleges. They are preceded by an introduction, in which the nature and use of logarithms are explained, and all necessary instruction given for using the tables. They are printed in large type with very open spacing. Compactness, simple arrangement, and figures large enough not to strain the eyes, are among the points in their favor.

Wentworth & Hill's Exercises in Arithmetic.

I. Exercise Manual. 12mo. Boards: vi + 282 pages. Mailing Price, 55 cents; for introduction, 50 cents. II. Examination Manual. 12mo. Boards. 148 pages. Mailing Price, 40 cents; Introduction Price, 35 cents. Both in one volume, 80 cents. Answers to both parts together, 10 cents.

THE first part (Exercise Manual) contains 3869 examples and problems for daily practice, classified and arranged in the common order; and the second part (Examination Manual) contains 300 examination-papers, progressive in character.

Wentworth & Hill's Exercises in Algebra.

I. Exercise Manual. 12mo. Boards. 232 pages. Mailing Price, 40 cents; Introduction Price, 35 cents. II. Examination Manual. 12mo. Boards. 159 pages. Mailing Price, 40 cents; Introduction Price, 35 cents. *Both in one volume*, 70 cents. *Answers to both parts together*, 25 cents.

THE first part (Exercise Manual) contains about 4500 problems classified and arranged according to the usual order of text-books in Algebra; and the second part (Examination Manual) contains nearly 300 examination-papers, progressive in character, and well adapted to cultivate skill and rapidity in solving problems.

<p>British Mail: All engaged in the practical work of education will appreciate these Manuals, as they are calculated to save the master</p>	<p>much precious time and labor, and to give his students the benefit of progressive and carefully thought-out exercises.</p>
---	---

Wentworth & Hill's Exercises in Geometry.

12mo. Cloth. 255 pages. Mailing Price, 80 cents; Introduction Price, 70 cents.

THE exercises consist of a great number of easy problems for beginners, and enough harder ones for more advanced pupils. The problems of each section are carefully graded, and some of the more difficult sections can be omitted without destroying the unity of the work. The book can be used in connection with any text-book on Geometry as soon as the geometrical processes of reasoning are well understood.

<p>Amelia W. Platter, High School, Indianapolis, Ind.: I find the subject so carefully graded, that I can</p>	<p>select propositions from it to supplement every stage of our work.</p>
--	---

Analytic Geometry.

By G. A. WENTWORTH. *Revised edition.* 12mo. Half morocco. xii + 301 pages. Mailing Price, \$1.35; for introduction, \$1.25; allowance in exchange, 30 cents.

THE aim of this work is to present the elementary parts of the subject in the best form for class-room use.

The connection between a locus and its equation is made perfectly clear in the opening chapter.

The exercises are well graded and designed to secure the best mental training.

By adding a supplement to each chapter provision is made for a shorter or more extended course, as the time given to the subject will permit.

The book is divided into chapters as follows :—

PART I. Plane Geometry. I. Loci and their Equations; II. The Straight Line; III. The Circle; IV. Different Systems of Co-ordinates; V. The Parabola; VI. The Ellipse; VII. The Hyperbola; VIII. Loci of the Second Order; IX. Higher Plane Curves.

PART II. Solid Geometry. I. The Point; II. The Plane; III. The Straight Line; IV. Surfaces of Revolution.

Dascom Greene, *Prof. of Mathematics and Astronomy, Rensselaer Polytechnic Institute, Troy, N.Y.*: It appears to be admirably adapted to the use of beginners, and is especially rich in examples for practical application of the principles of each chapter. The full and clear explanation of first principles given in the opening chapter is a new and highly commendable feature of the work. (Nov. 11, 1886.)

Geo. D. Olds, *Prof. of Mathematics, University of Rochester, N.Y.*: It is a most admirable little book. The author falls into line with what I believe to be the best modern tendency in text-books, — the avoidance of bulk and complexity. (Nov. 11, 1886.)

J. L. Patterson, *Teacher of Mathematics, Lawrenceville School, N.J.*: I do not know of any text-book for beginners in this subject which can compare with it for class-room use.

E. A. Paul, *Prin. of High School, Washington, D.C.*: I think it is to be commended for the same clearness of statement and simplicity of ar-

range-ment for which the author's other works are noted, and believe it to be especially adapted for advanced pupils, in high schools and academies, who wish to know something of the mysteries of loci and conic sections, and who have only a limited time for the work. (Nov. 12, 1886.)

Jos. J. Hardy, *Prof. of Mathematics, Lafayette College, Easton, Pa.*: The professor's experience has taught him what are the points which the boys find obscure, and he has generally been successful in devising a good way of elucidating these points. . . . Teachers will find it a very helpful manual. (Jan. 4, 1887.)

E. Miller, *Prof. of Mathematics, University of Kansas, Lawrence*: As a book for beginners, it is admirable in all its arrangements and features. The great number of problems scattered through it will largely relieve it of that abstract analysis which is so often a terror to students. The book is, like the other works of Professor Wentworth, a good thing. (Nov. 18, 1886.)

A Treatise on Plane Surveying.

By DANIEL CARHART, C.E., Professor of Civil Engineering in the Western University of Pennsylvania, Allegheny. Illustrated. 8vo. Half leather. xvii + 498 pages. Mailing Price, \$2.00; for introduction, \$1.80.

THIS work covers the whole ground of Plane Surveying. It illustrates and describes the instruments employed, their adjustments and uses; it exemplifies the best methods of solving the ordinary problems occurring in practice, and furnishes solutions for many special cases which not infrequently present themselves. It is the result of twenty years' experience in the field and technical schools, and the aim has been to make it extremely practical, having in mind always that to become a *reliable* surveyor the student needs frequently to manipulate the various surveying instruments in the field, to solve many examples in the class-room, and to exercise good judgment in all these operations. Not only, therefore, are the different methods of surveying treated, and directions for using the instruments given, but these are supplemented by various field exercises to be performed, by numerous examples to be wrought, and by many queries to be answered.

Chapter I. **Chain Surveying.**

" II. **Compass and Transit Surveying.**

" III. **Declination of the Needle.**

" IV. **Laying Out and Dividing Land.**

" V. **Plane Table Surveying.**

" VI. **Government Surveying.**

" VII. **City Surveying.** Including the Principles of Levelling.

" VIII. **Mine Surveying.** Including the Elements of Topography.

The following **Tables** have been added: Logarithms of numbers; Approximate equation of time; Logarithms of trigonometric functions; For determining with greater accuracy than the preceding; Lengths of degrees of latitude and longitude; Miscellaneous formulas, and equivalents of metres, chains, and feet; Traverse; Natural sines and cosines; Natural tangents and cotangents.

The judicial functions of surveyors, as given by Chief Justice Cooley, are set forth in an appendix.

The work is published just as this Catalogue goes to press, so that full notices cannot be given. Send for the special circular.

Byerly's Syllabi.

By W. E. BYERLY, Professor of Mathematics in Harvard University.
Each, 8 or 12 pages, 10 cents.

Syllabus of a Course in Plane Trigonometry.

Syllabus of a Course in Plane Analytical Geometry.

Syllabus of a Course in Plane Analytic Geometry. (Advanced Course.)

Syllabus of a Course in Analytical Geometry of Three Dimensions.

Syllabus of a Course on Modern Methods in Analytic Geometry.

Syllabus of a Course in the Theory of Equations.

Elements of the Differential and Integral Calculus.

With Examples and Applications. By J. M. TAYLOR, Professor of Mathematics in Madison University. 8vo. Cloth. 249 pages. Mailing Price, \$1.95; Introduction Price, \$1.80.

THE aim of this treatise is to present simply and concisely the fundamental problems of the Calculus, their solution, and more common applications. Its axiomatic datum is that the change of a variable, when not uniform, may be conceived as becoming uniform at any value of the variable.

It employs the conception of rates, which affords finite differentials, and also the simplest and most natural view of the problem of the Differential Calculus. This problem of *finding the relative rates of change of related variables* is afterwards reduced to that of finding the limit of the ratio of their simultaneous increments; and, in a final chapter, the latter problem is solved by the principles of infinitesimals.

Many theorems are proved both by the method of rates and that of limits, and thus each is made to throw light upon the other. The chapter on differentiation is followed by one on direct integration and its more important applications. Throughout the work there are numerous practical problems in Geometry and Mechanics, which serve to exhibit the power and use of the science, and to excite and keep alive the interest of the student.

The Nation, New York: It has two marked characteristics. In the first place, it is evidently a most carefully written book. There is nothing vague or slipshod in it. Nearly every sentence, certainly every theorem, seems to have been constructed with a strenuous effort to give it clearness and precision. This constant attention to the form of expression has enabled the author to be concise without becoming obscure. We are acquainted with no text-book of the calculus which compresses so much matter into so few pages, and at the same time leaves the impression that all that is necessary has been said. In the second place, the number of carefully selected examples, both of those worked out in full in illustration of the text, and of those left for the student to work out for himself, is extraordinary. From this point of view, those teachers and pupils who are accustomed to or prefer a different text-book, would still do well to provide themselves with this, regarding it merely as a collection of examples and without any reference to the text.

Elementary Co-ordinate Geometry.

By W. B. SMITH, Professor of Physics, Missouri State University. 12mo. Cloth. 312 pages. Mailing Price, \$2.15; for introduction, \$2.00.

WHILE in the study of Analytic Geometry either gain of knowledge or culture of mind may be sought, the latter object alone can justify placing it in a college curriculum. Yet the subject may be so pursued as to be of no great educational value. Mere calculation, or the solution of problems by algebraic processes, is a very inferior discipline of reason. Even geometry is not the best discipline. In all thinking, the real difficulty lies in forming clear notions of things. In doing this all the higher faculties are brought into play. It is this formation of concepts, therefore, that is the essential part of mental training. And it is in line with this idea that the present treatise has been composed. Professors of mathematics speak of it as the most exhaustive work on the subject yet issued in America; and in colleges where an easier text-book is required for the regular course, this will be found of great value for post-graduate study.

Wm. G. Peck, Prof. of Mathematics and Astronomy, Columbia College: I have read Dr. Smith's Co-ordinate Geometry from beginning to end with unflagging interest. Its well compacted pages contain an immense amount of matter, most admirably arranged. It is an excellent book, and the author is entitled to the thanks of every lover of mathematical science for this valuable contribution to its literature. I shall recommend its adoption as a text-book in our graduate course.

Academic Trigonometry : Plane and Spherical.

By T. M. BLAKSLÉE, Ph.D. (Yale), Professor of Mathematics in the University of Des Moines. 12mo. Paper. 33 pages. Mailing Price, 20 cents; for introduction, 15 cents.

THE Plane and Spherical portions are arranged on opposite pages.

The memory is aided by analogies, and it is believed that the entire subject can be mastered in less time than is usually given to Plane Trigonometry alone, as the work contains but 29 pages of text.

The Plane portion is compact, and complete in itself.

Examples of Differential Equations.

By GEORGE A. OSBORNE, Professor of Mathematics in the Massachusetts Institute of Technology, Boston. 12mo. Cloth. vii + 50 pages. Mailing Price, 60 cents; for introduction, 50 cents.

A SERIES of nearly three hundred examples with answers, systematically arranged and grouped under the different cases, and accompanied by concise rules for the solution of each case.

Selden J. Coffin, lately Prof. of *Mathematics*, Lafayette College : Its appearance is most timely, and it supplies a manifest want.

Determinants.

The Theory of Determinants: an Elementary Treatise. By PAUL H. HANUS, B.S., recently Professor of Mathematics in the University of Colorado, now Principal of West High School, Denver, Col. 8vo. Cloth. viii + 217 pages. Mailing Price, \$1.90; for introduction, \$1.80.

THIS book is written especially for those who have had no previous knowledge of the subject, and is therefore adapted to self-instruction as well as to the needs of the class-room. The subject is at first presented in a very simple manner. As the reader advances, less and less attention is given to details. Throughout the entire work it is the constant aim to arouse and enliven the reader's interest, by first showing how the various concepts have arisen naturally, and by giving such applications as can be presented without exceeding the limits of the treatise. The work is sufficiently comprehensive to enable the student who has mastered the volume to use the determinant notation with ease, and to pursue his further reading in the modern higher algebra with pleasure and profit.

GENERAL LIBRARY
UNIVERSITY OF CALIFORNIA—BERKELEY

RETURN TO DESK FROM WHICH BORROWED

This book is due on the last date stamped below, or on the
date to which renewed.

Renewed books are subject to immediate recall.

18 Apr '55 KW

21 1956 11

22 Oct '56 WJ

REC'D LD

OCT 8 1956

12 Mar '63 WS

REC'D LD

FEB 26 1963

YB 56843

797961

QA531

B6

UNIVERSITY OF CALIFORNIA LIBRARY

